

fragments in the untranslated portion of the mRNA encoding the antibody or antibody fragment. The 3' untranslated regions also include transcription termination sites.

Suitable vectors containing one or more of the above listed components and the desired coding and control sequences are constructed by standard ligation techniques. Isolated plasmids or DNA fragments are cleaved, tailored, and religated in the form desired to generate the plasmids required.

For analysis to confirm correct sequences in plasmids constructed, the ligation mixtures are used to transform *E. coli* K12 strain 294 (ATCC 31,446) and successful transformants selected by ampicillin or tetracycline resistance where appropriate. Plasmids from the transformants are prepared, analyzed by restriction endonuclease digestion, and/or sequenced by the method of Messing *et al.*, Nucleic Acids Res., 9: 309 (1981) or by the method of Maxam *et al.*, Methods in Enzymology, 65: 499 (1980).

Particularly useful in the practice of this invention are expression vectors that provide for the transient expression in mammalian cells of DNA encoding the antibody or antibody fragment. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of a desired polypeptide encoded by the expression vector.

Other methods, vectors, and host cells suitable for adaptation to the synthesis of the antibody or antibody fragment in recombinant vertebrate cell culture are described in Gething *et al.*, Nature, 293: 620-625 (1981); Mantei *et al.*, Nature, 281: 40-46 (1979); Levinson *et al.*, EP 117,060; and EP 117,058. A particularly useful plasmid for mammalian cell culture expression of the IgE peptide antagonist is pRK5 (EP pub. no. 307,247) or pSVI6B (PCT pub. no. WO 91/08291 published 13 June 1991).

### C. Selection and Transformation of Host Cells

Suitable host cells for cloning or expressing the vectors herein are the prokaryote, yeast, or higher eukaryote cells described above. Suitable prokaryotes include eubacteria, such as Gram-negative or Gram-positive organisms, for example, *E. coli*, *Bacilli* such as *B. subtilis*,